

## **Facility to Facility Transport Criteria – During Transport**

|   | BLS Transport | ALS Transport | CCT Transport |
|---|---------------|---------------|---------------|
| 02  | х             |               |               |
| IV Lock   | x             |               |               |
| Tracheostomy Management – Suctioning                          | х             |               |               |
| Patient PCA   | х             |               |               |
| Cardiac Monitoring  |               | Х             |               |
| Pacing – Transcutaneous                                       |               | Х             |               |
| CPAP only – Non Ventilator                                    |               | Х             |               |
| IV Fluid Infusion (NS, D5W, LR - No medication being infused) |               | Х             |               |
| ACLS Medications  |               | X             |               |
| Pain Management with IVP Medication                           |               | X             |               |
| PIC or Shunt Monitoring                                       |               | X             |               |
| BiPAP   |               |               | x             |
| Ventilator  |               |               | x             |
| Volume Controlled Medications - Med Pump                      |               |               | x             |
| Continuous Infusion w/ PCA                                    |               |               | X             |
| Blood products  |               |               | x             |
| Chest Tube Monitoring   |               |               | x             |
| IV Programmable Volume Device                                 |               |               | X             |
| Arterial Line – Monitoring and Access                         |               |               | Х             |
| Central Line  |               |               | X             |

TransportCriteria/KCEMSS 4/12/2024

